

AD-A044 817

MARYLAND UNIV COLLEGE PARK DEPT OF COMPUTER SCIENCE F/G 12/1  
SENSITIVITY COEFFICIENTS FOR THE EFFECTS OF ERRORS IN THE INDEP--ETC(U)  
SEP 77 G W STEWART N00014-76-C-0391

UNCLASSIFIED


TR-571

NL

1 OF 1  
ADA  
044817



END  
DATE  
FILMED  
10-77  
DDC



NATIONAL BUREAU OF STANDARDS  
MICROCOPY RESOLUTION TEST CHART

AD A 044817

12  
B.S.



COMPUTER SCIENCE  
TECHNICAL REPORT SERIES

DDC  
100  
1977  
C



UNIVERSITY OF MARYLAND  
COLLEGE PARK, MARYLAND

20742

AD No. \_\_\_\_\_  
DDC FILE COPY.

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

14 TR-571

12 10 p.

9 Technical Report TR-571  
ONR-N00014-76-C-0391-571

11 Sep 1977

15 6 Sensitivity Coefficients  
for the Effects of Errors in  
the Independent Variables in  
a Linear Regression.

by

10 G. W. Stewart



#### Abstract

This paper is concerned with errors in the observed values of the independent variables of a linear regression. We propose sensitivity coefficients to measure the effects of these errors and show that they can easily be computed from quantities ordinarily calculated in performing the regression.

This work was supported in part by the Office of Naval Research under Contract No. N00014-76-C-0391.

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

409 022

Sensitivity Coefficients  
for the Effects of Errors in  
the Independent Variables in  
a Linear Regression

G. W. Stewart

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED JUSTIFICATION	
DISTRIBUTION/AVAILABILITY CODES	
SPECIAL	
A	

1. Introduction

In this paper we shall be concerned with the regression problem

$$\text{minimize } \|y - \tilde{X}\beta\|^2,$$

where  $\tilde{X}$  is an  $n \times p$  matrix of rank  $p$ ,  $y$  is an  $n$ -vector, and  $\|\cdot\|$  denotes the usual Euclidean vector norm.\* The problem has the unique solution

$$(1.1) \quad \beta = \tilde{X}^+ y$$

where  $\tilde{X}^+ = (\tilde{X}'\tilde{X})^{-1}\tilde{X}'$  is the pseudo-inverse of  $\tilde{X}$ .

Although classical regression theory concerns itself with the statistical analysis of errors in the vector  $y$ , it frequently happens that the design matrix  $\tilde{X}$  is itself contaminated with errors, so that one is effectively working with a perturbed matrix  $\tilde{X} + E$ . For example, the columns of  $\tilde{X}$  may be measured by means of some instrument for which the originator of the problem can only give crude error estimates.

In this case the data analyst is faced with the problem of deciding when the effects of the errors can be ignored. The problem is especially

\* In the sequel  $\|\cdot\|$  will also denote the spectral matrix norm defined by  $\|A\| = \sup \{\|Ax\| : \|x\|=1\}$ . See [3] for details.

critical in general purpose regression routines, where it is desirable to provide the user with a set of easily interpretable numbers that indicate the magnitude of the effects of the errors.

A partial solution is provided by the perturbation theory for the regression problem (for a survey of this theory see [4]). This theory bounds perturbations in  $\beta$  in terms of  $\|E\|$  and of the "condition number"  $\kappa = \|\underline{X}\| \|\underline{X}^\dagger\|$ . Although the results of this theory shed considerable light on the behavior of regression problems under perturbations in  $\underline{X}$ , they are unsatisfactory in practice for two reasons. First they bound only the norm of the perturbation in  $\beta$ , so that a large perturbation in one component can conceal the fact that the others have small perturbations. More important, they are not scale invariant; changing the scale of the columns of  $\underline{X}$  will change  $\kappa$ , even though the statistical problem is essentially unaltered. This phenomenon makes the results quite difficult to interpret.

Taking a different approach, Beaton, Rubin, and Barone [1] have derived measures of sensitivity that to some extent answer the above objections. However, it is assumed that the errors are unbiased and  $n$  is large. Swindel and Bauer [5] have derived a useful bound for the relative bias in  $\beta$ , which measures the relative effects of perturbations in  $\underline{X}$  compared to the usual statistical errors in  $\underline{X}$ . In this paper we shall derive coefficients  $\gamma_{ij}$  that measure the sensitivity of  $\beta_i$  to changes in column  $j$  of  $\underline{X}$ . Specifically  $\gamma_{ij}$  is the norm of the Frechet



derivative of  $\beta_i$  regarded as a function of the  $j$ -th column of  $\tilde{X}$ . If  $\epsilon$  is the norm of the perturbation of the  $i$ -th column of  $\tilde{X}$ , then  $\gamma_{ij}\epsilon$  will be an asymptotic bound on the perturbation induced in  $\beta_j$ .\*

## 2. Derivation of the Coefficients

Although it is in principle possible to calculate the required derivatives directly from the normal equations  $(\tilde{X}'\tilde{X})\beta = \tilde{X}'y$ , we prefer to approach the problem through a first order perturbation theorem that is useful in its own right.

Theorem 2.1. In the notation of the last section, let  $\beta$  be the solution of the regression problem (1.1), and let  $\tilde{r} = y - \tilde{X}\beta$  be the corresponding residual vector. Let  $\tilde{E}$  be an  $n \times p$  matrix. If

$$(2.1) \quad \|\tilde{X}^+ \tilde{E}\| < 1,$$

then  $\tilde{X} + \tilde{E}$  has rank  $p$  so that there is a unique solution  $\tilde{\beta}$  of the problem

$$\text{minimize } \|y - (\tilde{X} + \tilde{E})\tilde{\beta}\|^2.$$

Moreover, as  $\tilde{E}$  approaches zero

$$\tilde{\beta} = \beta - \tilde{X} \tilde{E} \beta + (\tilde{X}'\tilde{X})^{-1} \tilde{E}' \tilde{r} + O(\|\tilde{E}\|^2).$$

Proof. For a proof that (2.1) implies that  $\tilde{X} + \tilde{E}$  has full rank, see [4]. This implies that for all sufficiently small  $\tilde{E}$ ,  $\tilde{\beta}$  exists and is given by

---

\* Here, and throughout this note, the term asymptotic refers to behavior for small  $\epsilon$ , not large  $n$ .

$$\tilde{\beta} = [(X+E)'(X+E)]^{-1}(X+E)\gamma.$$

Now it is well known [3] that if  $\tilde{P}$  is sufficiently small, then  $\tilde{I} + \tilde{P}$  is nonsingular and

$$(\tilde{I} + \tilde{P})^{-1} = \tilde{I} - \tilde{P} + O(\|\tilde{P}\|^2).$$

It follows that

$$\begin{aligned} [(X+E)'(X+E)]^{-1} &= (X'X + X'E + E'X + E'E)^{-1} \\ &= \{X'X[I + (X'X)^{-1}(X'E + E'X + E'E)]\}^{-1} \\ &= [I - (X'X)^{-1}(X'E + E'X)](X'X)^{-1} + O(\|E\|^2). \end{aligned}$$

Hence

$$\begin{aligned} \tilde{\beta} &= (X'X)^{-1}X'\gamma + (X'X)^{-1}E'\gamma - (X'X)^{-1}X'E(X'X)^{-1}X'\gamma \\ &\quad - (X'X)^{-1}E'X(X'X)^{-1}X'\gamma + O(\|E\|^2) \\ &= \beta - X^+E\gamma + (X'X)^{-1}E'(\gamma - X\beta) \\ &= \beta - X^+E\gamma + (X'X)^{-1}E^T r. \quad \square \end{aligned}$$

Theorem 2.1 immediately gives an expression for the Frechet derivative of  $\beta_i$  regarded as a function of the  $j$ -th column of  $X$ .

Corollary 2.2. Let  $\beta_i = f_{ij}(x_j)$ , where  $x_j$  denotes the  $j$ -th column of  $X$ . Then

$$(2.2) \quad Df_{ij} = -\beta_j e_i^T X^+ + e_i' (X'X)^{-1} e_j r^T.$$



Proof. The Frechet derivative  $\tilde{Df}_{ij}$  is the unique row vector satisfying

$$(2.3) \quad f_{ij}(\tilde{x}_j + \tilde{e}) = \beta_i + \tilde{Df}_{ij}\tilde{e} + O(\|\tilde{e}\|^2).$$

Let  $\tilde{e}_j$  denote the  $j$ -th unit vector. Then a perturbation  $\tilde{e}$  in  $\tilde{x}_j$  amounts to adding to  $\tilde{X}$  the matrix  $\tilde{E} = \tilde{e}\tilde{e}_j^T$ . Hence from Theorem 2.1

$$\begin{aligned} f_{ij}(\tilde{x}_j + \tilde{e}) &= \tilde{e}_i'[\beta - \tilde{X}(\tilde{e}\tilde{e}_j')\beta + (\tilde{X}'\tilde{X})^{-1}(\tilde{e}\tilde{e}_j')'r] + O(\|\tilde{e}\|^2) \\ &= \beta_i - \beta_j \tilde{e}_i' \tilde{X} \tilde{e} + \tilde{e}_i' (\tilde{X}\tilde{X})^{-1} \tilde{e}_j \tilde{e}' r + O(\|\tilde{e}\|^2) \\ &= \beta_i - (\beta_j \tilde{e}_i' \tilde{X} - \tilde{e}_i' (\tilde{X}'\tilde{X})^{-1} \tilde{e}_j r') \tilde{e} + O(\|\tilde{e}\|^2), \end{aligned}$$

which shows that  $f'_{ij}$  defined by (2.2) satisfies (2.3).  $\square$

Corollary 2.3. Let  $C = (\tilde{X}'\tilde{X})^{-1}$ . Then

$$\|\tilde{Df}_{ij}\| = \gamma_{ij} = \sqrt{|\beta_j|^2 c_{ii} + \|r\|^2 c_{ij}^2}.$$

Proof. The vector  $r$  is orthogonal to the column space of  $\tilde{X}$ , which is the same as the row space of  $\tilde{X}^T$ . Hence  $\tilde{e}_i^T \tilde{X}^T$  and  $r'$  are orthogonal, and

$$\|\tilde{Df}_{ij}\|^2 = |\beta_j|^2 \|\tilde{e}_i \tilde{X}^T\|^2 + c_{ij}^2 \|r\|^2.$$

The proof will be complete if we can show that  $\|\tilde{e}_i \tilde{X}^T\|^2 = c_{ii}$ . But

$$\|\tilde{e}_i \tilde{X}^T\|^2 = \|\tilde{e}_i' (\tilde{X}'\tilde{X})^{-1} \tilde{X}'\|^2 = \tilde{e}_i' (\tilde{X}'\tilde{X})^{-1} \tilde{X}' \tilde{X} (\tilde{X}'\tilde{X})^{-1} \tilde{e}_i = \tilde{e}_i' (\tilde{X}'\tilde{X})^{-1} \tilde{e}_i = c_{ii}. \quad \square$$

### 3. Applications

It should be observed that the numbers  $\gamma_{ij}$  can be calculated from the quantities that are usually computed in the course of the regression. For example, if sweep techniques (e.g. see [2]) have been used to solve the problem, the numbers  $c_{ij}$  will be available, as will  $\|\tilde{r}\|^2$ , since it is simply the residual sum of squares.

The results are readily interpretable. If a bound  $\epsilon_j$  on the norm of the error in the  $j$ -th column is available, then the error  $\delta_i$  induced in  $\beta_i$  is asymptotically bounded by  $\gamma_{ij}\epsilon_j$ . If this number is too large, the problem requires further study. In interpreting the results it should always be borne in mind that  $\gamma_{ij}\epsilon_j$  is a first order bound. A large value is a signal that something may be wrong, but if  $\epsilon_j$  is so large that the first order approximation is not applicable, then the difficulties may turn out to be illusory.

A particularly attractive feature of the asymptotic bounds is that, since they deal with individual components of  $\beta$ , their interpretation is independent of the scaling of the columns of  $\tilde{X}$ . This is particularly apparent if the bounds are cast in terms of relative error in the form

$$\frac{|\delta_i|}{|\beta_i|} \lesssim \left( \frac{\gamma_{ij} \|\tilde{x}_j\|}{\|\beta_i\|} \right) \frac{\epsilon_j}{\|\tilde{x}_j\|}.$$

Now

$$(3.1) \quad \frac{\gamma_{ij} \|\tilde{x}_j\|}{|\beta_i|} = \left[ (|\beta_j|^2 \|\tilde{x}_j\|^2) \left( \frac{c_{ii}}{|\beta_i|^2} \right) + (\|\tilde{r}\|^2) \left( \frac{|c_{ij}|^2 \|\tilde{x}_j\|^2}{|\beta_i|^2} \right) \right]^{1/2},$$

and each parenthesized term in the right hand side of (3.1) is easily seen to be invariant under scaling of the columns of  $\tilde{X}$ . Indeed, in some applications where the  $\beta_i$ 's are known to be bounded away from zero it may be more appropriate to report  $\gamma_{ij} \|\tilde{x}_j\|/|\beta_i|$  than  $\gamma_{ij}$ .

In deriving our bounds, we have used the Cauchy-Schwarz inequality  $\|\tilde{x}^*y\| \leq \|\tilde{x}\|\|y\|$ , an inequality which is usually pessimistic, since it must account for the worst case where  $\tilde{x}$  and  $y$  are dependent. If we are willing to assume more about the perturbation  $e$  in  $\tilde{x}_j$ , then we may be able to say more. For example, we have the following consequence of Corollary 2.3.

**Corollary 3.1.** Let  $e \in N(0, \sigma^2 I)$ . Then  $Df_{ij}e$  is normally distributed with mean zero and standard deviation  $\gamma_{ij}\sigma$ .

#### References

1. A. E. Beaton, D. B. Rubin, and J. L. Barne, The acceptability of regression solutions: another look at computational accuracy, J. Amer. Stat. Assoc. 71 (1976) 158-168.
2. A. P. Dempster, Elements of Continuous Multivariate Analysis, Addison-Wesley, Reading, Massachusetts (1969).
3. G. W. Stewart, Introduction to Matrix Computations, Academic Press, New York (1973).
4. \_\_\_\_\_, On the perturbation of pseudo-inverses, projections, and linear least squares problems, to appear SIAM Rev.
5. B. F. Swindel and D. R. Bower, Rounding errors in the independent variables in a general linear model, Technometrics 14 (1972) 215-218.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ONR-N00014-76-C-0391-571	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) SENSITIVITY COEFFICIENTS FOR THE EFFECTS OF ERRORS IN THE INDEPENDENT VARIABLES IN A LINEAR REGRESSION		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) G. W. Stewart		6. PERFORMING ORG. REPORT NUMBER TR-571 ✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Computer Science ✓ University of Maryland College Park, MD 20742		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0391 ✓
11. CONTROLLING OFFICE NAME AND ADDRESS Mathematics Branch Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE September 1977
		13. NUMBER OF PAGES 9
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) regression least squares errors in variables sensitivity		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This paper is concerned with errors in the observed values of the independent variables of a linear regression. We propose sensitivity coefficients to measure the effects of these errors and show that they can easily be computed from quantities ordinarily calculated in performing the regression.		